

付表-C.1.7 長周期波高の出現分布（酒田）

酒田 冬季：'07/12～'07/2									
波高階級 (cm)	4√m0L30-60	4√m0L60-300	4√m0L30-	観測数	出現率	観測数	出現率	観測数	出現率
- 1	0	0.0	0	0	0.0	0	0.0	0	0.0
1 - 5	691	63.3	357	32.7	53	4.9			
5 - 10	350	32.1	480	44.0	616	56.4			
10 - 20	39	3.6	212	19.4	341	31.2			
20 - 50	5	0.5	38	3.5	71	6.5			
50 -	2	0.2	0	0.0	6	0.5			
欠測					5	0.5			
全観測				1092					
平均値	5.0		7.8		10.7				

酒田 春季：'08/3～'08/5									
波高階級 (cm)	4√m0L30-60	4√m0L60-300	4√m0L30-	観測数	出現率	観測数	出現率	観測数	出現率
- 1	0	0.0	0	0.0	0	0.0	0	0.0	0.0
1 - 5	1016	92.0	746	67.6	293	26.5			
5 - 10	72	6.5	320	29.0	712	64.5			
10 - 20	0	0.0	22	2.0	83	7.5			
20 - 50	0	0.0	0	0.0	0	0.0			
50 -	0	0.0	0	0.0	0	0.0			
欠測					16	1.4			
全観測				1104					
平均値	3.1		4.5		6.5				

酒田 夏季：'08/6～'08/8									
波高階級 (cm)	4√m0L30-60	4√m0L60-300	4√m0L30-	観測数	出現率	観測数	出現率	観測数	出現率
- 1	2	0.2	0	0.0	0	0.0			
1 - 5	1068	96.7	959	86.9	621	56.2			
5 - 10	1	0.1	112	10.1	448	40.6			
10 - 20	0	0.0	0	0.0	2	0.2			
20 - 50	0	0.0	0	0.0	0	0.0			
50 -	0	0.0	0	0.0	0	0.0			
欠測					33	3.0			
全観測				1104					
平均値	2.3		3.2		4.9				

酒田 秋季：'08/9～'08/11									
波高階級 (cm)	4√m0L30-60	4√m0L60-300	4√m0L30-	観測数	出現率	観測数	出現率	観測数	出現率
- 1	0	0.0	0	0.0	0	0.0			
1 - 5	863	79.0	645	59.1	351	32.1			
5 - 10	172	15.8	284	26.0	474	43.4			
10 - 20	3	0.3	109	10.0	191	17.5			
20 - 50	0	0.0	0	0.0	22	2.0			
50 -	0	0.0	0	0.0	0	0.0			
欠測					54	4.9			
全観測				1092					
平均値	3.4		5.4		7.5				

酒田 通年：'08/1～'08/12									
波高階級 (cm)	4√m0L30-60	4√m0L60-300	4√m0L30-	観測数	出現率	観測数	出現率	観測数	出現率
- 1	2	0.0	0	0.0	0	0.0			
1 - 5	3614	82.3	2677	61.0	1304	29.7			
5 - 10	584	13.3	1200	27.3	2227	50.7			
10 - 20	78	1.8	327	7.4	612	13.9			
20 - 50	5	0.1	81	1.8	136	3.1			
50 -	2	0.0	0	0.0	6	0.1			
欠測					107	2.4			
全観測				4392					
平均値	3.5		5.4		7.7				